

Survival Ensembles

Torsten Hothorn^{1,*}, Peter Bühlmann², Sandrine Dudoit³,
Annette Molinaro⁴ and Mark J. van der Laan³

¹Institut für Statistik
Ludwig-Maximilians-Universität München
Ludwigstraße 33, D-80539 München, Germany
Tel: ++49-9131-8522707
Fax: ++49-9131-8525740
`Torsten.Hothorn@R-project.org`

²Seminar für Statistik, ETH Zürich, CH-8032 Zürich, Switzerland
`buehlmann@stat.math.ethz.ch`

³Division of Biostatistics, University of California, Berkeley
140 Earl Warren Hall, #7360, Berkeley, CA 94720-7360, USA
`sandrine@stat.Berkeley.EDU`
`laan@stat.Berkeley.EDU`

⁴Division of Biostatistics, Epidemiology and Public Health
Yale University School of Medicine, 206 LEPH
60 College Street PO Box 208034, New Haven CT 06520-8034
`annette.molinaro@yale.edu`

1 Illustrations and Applications

This document reproduces the data analyses presented in [Hothorn et al. \(2006\)](#). For a description of the theory behind applications shown here we refer to the original manuscript. The results differ slightly due to technical changes or bug-fixes in **mboost** that have been implemented after the paper was printed.

1.1 Acute myeloid leukemia

Data preprocessing Compute IPC weights, define risk score and set up learning sample:

```
R> AMLw <- IPCweights(Surv(clinical$time, clinical$event))  
R> risk <- rep(0, nrow(clinical))
```

```

R> rlev <- levels(clinical[, "Cytogenetic.group"])
R> risk[clinical[, "Cytogenetic.group"] %in% rlev[c(7,
8, 4)]] <- "low"
R> risk[clinical[, "Cytogenetic.group"] %in% rlev[c(5,
9)]] <- "intermediate"
R> risk[clinical[, "Cytogenetic.group"] %in% rlev[-c(4,
5, 7, 8, 9)]] <- "high"
R> risk <- as.factor(risk)
R> AMLlearn <- cbind(clinical[, c("time", "Sex",
"Age", "LDH", "WBC", "FLT3.aberration.", "MLL.PTD",
"Tx.Group.")], risk = risk, iexpressions[,
colnames(iexpressions) %in% selgenes[["Clone.ID"]]])
R> cc <- complete.cases(AMLlearn)
R> AMLlearn <- AMLlearn[AMLw > 0 & cc, ]
R> AMLw <- AMLw[AMLw > 0 & cc]

```

Model fitting Fit random forest for censored data

```

R> ctrl <- cforest_control(mincriterion = 0.1, mtry = 5,
minsplitt = 5, ntree = 250)
R> AMLrf <- cforest(I(log(time)) ~ ., data = AMLlearn,
control = ctrl, weights = AMLw)

```

and L_2 Boosting for censored data

```

R> AML12b <- glmboost(I(log(time)) ~ ., data = AMLlearn,
weights = AMLw, control = boost_control(mstop = 5000))

```

Compute fitted values

```

R> AML12b <- AML12b[mstop(aic)]
R> cAML <- coef(AML12b)
R> cAML[abs(cAML) > 0]

```

(Intercept)	Age	WBC
0.03094981	0.00854937	-0.00364371
MLL.PTDyes	Tx.Group.AUTO	Tx.Group.IC
-0.50709786	0.90185340	0.04037578
Tx.Group.Ind	riskintermediate	`IMAGE:145643`
-1.86134842	0.11825619	0.19788355
`IMAGE:2542486`	`IMAGE:345601`	`IMAGE:377560`
0.00442375	0.02935101	0.11000322
`IMAGE:428782`	`IMAGE:2043415`	`IMAGE:1584563`
0.01010658	0.05911671	-0.17883619
`IMAGE:347035`	`IMAGE:262695`	`IMAGE:950479`
-0.03307600	0.00080156	0.09049309
`IMAGE:898305`	`IMAGE:1472689`	`IMAGE:150702`
0.00523016	0.03498572	0.01367553
`IMAGE:1526826`	`IMAGE:66507`	`IMAGE:786302`
-0.01805326	0.00399127	0.08941300
`IMAGE:243614`	`IMAGE:417884`	`IMAGE:1592006`

```
R> plot(aic <- AIC(AML12b))
```




Figure 1: AIC criterion for AML data.

```

-0.05776062      -0.04890054      -0.02269622
`IMAGE:1917063`  `IMAGE:884333`  `IMAGE:133273`
-0.06536720      0.04189990      0.06594787
`IMAGE:950888`  `IMAGE:809533`  `IMAGE:49389`
0.02027810      -0.15986981      0.06352703
`IMAGE:789357`  `IMAGE:142139`  `IMAGE:1558053`
-0.01252187      0.00089307      0.07795515
`IMAGE:856174`  `IMAGE:504421`  `IMAGE:435036`
0.01115234      0.06861766      0.06094620
`IMAGE:491751`  `IMAGE:782835`  `IMAGE:52930`
0.04336285      -0.17924185      -0.03503330
`IMAGE:2545705`  `IMAGE:756405`  `IMAGE:502664`
-0.09886616      0.07713650      0.03620466
`IMAGE:129032`  `IMAGE:1610168`  `IMAGE:327676`
-0.31322459      0.01260374      -0.02117310
`IMAGE:69002`  `IMAGE:121551`  `IMAGE:2019101`
-0.41671336      -0.08107446      -0.06531175
`IMAGE:1456160`  `IMAGE:430318`  `IMAGE:2566064`
-0.10208684      -0.07297586      0.06126683
`IMAGE:74537`  `IMAGE:1606557`  `IMAGE:306812`
0.04523784      0.14243526      0.03504441
`IMAGE:565083`  `IMAGE:843028`  `IMAGE:68794`
0.29555347      0.05619983      0.23722775
`IMAGE:488505`  `IMAGE:167205`  `IMAGE:291756`
0.33464829      0.00217136      0.04973319
`IMAGE:810801`  `IMAGE:1702742`  `IMAGE:380462`
0.08725523      -0.04428190      -0.13182519
`IMAGE:154472`  `IMAGE:302540`  `IMAGE:135221`
-0.24723347      0.17175129      -0.01972168
`IMAGE:1567220`  `IMAGE:594630`
0.02473376      -0.07396882

```

```

R> AMLprf <- predict(AMLrfr, newdata = AMLlearn)
R> AMLpb <- predict(AMLl2b, newdata = AMLlearn)

```

1.2 Node-positive breast cancer

Data preprocessing Compute IPC weights and set up learning sample:

```

R> data("GBSG2", package = "ipred")
R> GBSG2w <- IPCweights(Surv(GBSG2$time, GBSG2$cens))
R> GBSG2learn <- cbind(GBSG2[, -which(names(GBSG2) %in%
      c("time", "cens"))], ltime = log(GBSG2$time))
R> n <- nrow(GBSG2learn)

```

Model fitting

```

R> LMmod <- lm(ltime ~ ., data = GBSG2learn, weights = GBSG2w)
R> LMerisk <- sum((GBSG2learn$ltime - predict(LMmod))^2 *
      GBSG2w)/n
R> TRmod <- rpart(ltime ~ ., data = GBSG2learn, weights = GBSG2w)
R> TRerisk <- sum((GBSG2learn$ltime - predict(TRmod))^2 *

```




Figure 2: AML data: Reproduction of Figure 1.

```

GBSG2w)/n
R> ctrl <- cforest_control(mincriterion = qnorm(0.95),
  mtry = 5, minsplit = 5, ntree = 100)
R> RFmod <- cforest(ltime ~ ., data = GBSG2learn,
  weights = GBSG2w, control = ctrl)
R> L2Bmod <- glmboost(ltime ~ ., data = GBSG2learn,
  weights = GBSG2w, control = boost_control(mstop = 250))
R> L2BHubermod <- glmboost(ltime ~ ., data = GBSG2learn,
  weights = GBSG2w, family = Huber(d = log(2)))

```

Compute fitted values:

```

R> GBSG2Hp <- predict(L2BHubermod, newdata = GBSG2learn)
R> L2Berisk <- sum((GBSG2learn$ltime - predict(L2Bmod,
  newdata = GBSG2learn))^2 * GBSG2w)/n
R> RFerisk <- sum((GBSG2learn$ltime - predict(RFmod,
  newdata = GBSG2learn))^2 * GBSG2w)/n

```

```
R> plot(aic <- AIC(L2Bmod))
```




Figure 3: AIC criterion for GBSG2 data.




Figure 4: GBSG-2 data: Reproduction of Figure 3.




Figure 5: GBSG-2 data: Reproduction of Figure 5.




Figure 6: GBSG-2 data: Reproduction of Figure 6.




Figure 7: GBSG-2 data: Reproduction of Figure 7.

References

- T. Hothorn, P. Bühlmann, S. Dudoit, A. Molinaro, and M. van der Laan. Survival ensembles. *Biostatistics*, 7:355–373, 2006.