

Package ‘demofit’

August 20, 2026

Title Parametric Mortality Curve Fitting and Mortality Forecasting Tools

Version 0.1.5

Author Jackie Li [aut, cre, cph]

Maintainer Jackie Li <jackieli@smu.edu.sg>

Description Provides tools for fitting parametric mortality curves. Implements multiple optimisation strategies to enhance robustness and stability of parameter estimation. Offers tools for forecasting mortality rates guided by mortality curves.

For modelling details see:

Tabeau (2001) <[doi:10.1007/0-306-47562-6_1](https://doi.org/10.1007/0-306-47562-6_1)>,

Renshaw and Haberman (2006) <[doi:10.1016/j.insmatheco.2005.12.001](https://doi.org/10.1016/j.insmatheco.2005.12.001)>,

Cairns et al. (2009) <[doi:10.1080/10920277.2009.10597538](https://doi.org/10.1080/10920277.2009.10597538)>,

Li and Lee (2005) <[doi:10.1353/dem.2005.0021](https://doi.org/10.1353/dem.2005.0021)>.

License GPL-3

Encoding UTF-8

RoxygenNote 7.3.3

Imports forecast, minpack.lm, MortalityLaws, NlcOptim, stats, MASS, graphics, grDevices

Suggests knitr, rmarkdown

VignetteBuilder knitr

NeedsCompilation no

Repository CRAN

Date/Publication 2026-08-19 22:20:23 UTC

Contents

APCS	2
CAES	4
CBDCS	5
CBDQCS	6
CBDS	8

CC	9
CFM2S	10
CFMS	11
curveS3	13
ENI	13
ENS	15
FCS	16
LCLS	17
LCS	19
MC	20
modelS3	22
mortalitymatrix	23
PAPCS	24
PCAES	25
PCBDCS	27
PCBDQCS	28
PCBDS	29
PCFM2S	31
PCFMS	32
PFCS	34
PLCS	36
PRHS	37
RHS	39
STARS	40
Index	43

APCS

*Age-period-cohort model***Description**

Fits and forecasts mortality rates using age-period-cohort model.

Usage

```
APCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2,

where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).

h forecast horizon (default = 10).

jumpoff if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The age-period-cohort (APC) model is specified as

$$\ln(m_{x,t}) = \alpha_x + \kappa_t + \gamma_{t-x} + \epsilon_{x,t}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c . Constraints include sum of κ_t is zero and and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class APCS with associated S3 methods `coef`, `forecast` (which = 1 for smoothed (default); which = 2 for raw), `plot`, `residuals`, and `simulate` (nsim for setting number of simulations; seed for initialising random number generator).

References

Bray, I. (2002). Application of Markov chain Monte Carlo methods to projecting cancer incidence and mortality. *Journal of the Royal Statistical Society Series C*, 51(2), 151-164.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- rep(1/30,30)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- APCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CAES

*Common age effect model***Description**

Fits and forecasts mortality rates of two populations using common age effect model.

Usage

```
CAES(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common age effect (CAE) model is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + \beta_x \kappa_{t,i} + \epsilon_{x,t,i}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{t,i}$. Constraints include sum of β_x is one and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CAES with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Kleinow, T. (2015). A common age effect model for the mortality of multiple populations. Insurance: Mathematics and Economics, 63(C), 147-152.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
b <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
k1 <- c(8.68,8.34,7.99,6.87,8.18,5.73,4.83,5.20,2.74,3.22,
2.99,1.59,1.67,-0.65,-0.39,-1.07,-0.95,-2.78,-3.46,-2.45,
-4.12,-4.66,-4.98,-4.58,-6.30,-4.39,-5.56,-6.52,-8.26,-6.92)
k2 <- c(11.81,11.01,10.59,10.40,9.75,8.15,6.07,6.45,4.60,4.57,
4.15,1.49,1.77,-1.08,-1.44,-0.96,-1.66,-2.25,-4.67,-4.62,
-4.38,-6.37,-6.27,-6.91,-8.22,-7.35,-8.39,-7.87,-9.72,-8.65)
set.seed(123)
M1 <- exp(outer(k1,b)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CAES(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDCS

CBD with cohort model

Description

Fits and forecasts mortality rates using CBD with cohort model.

Usage

```
CBDCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).

jumpoff if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with cohort (M6) model is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \gamma_{t-x} + \epsilon_{x,t}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, and γ_c . Constraints include sum of γ_c is zero and sum of $c\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class CBDCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDQCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDQCS

CBD with curvature and cohort model

Description

Fits and forecasts mortality rates using CBD with curvature and cohort model.

Usage

```
CBDQCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with curvature and cohort (M7) model is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \kappa_{3,t}((x - \bar{x})^2 - \sigma^2) + \gamma_{t-x} + \epsilon_{x,t}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, $\kappa_{3,t}$, and γ_c . Constraints include sum of γ_c is zero, sum of $c\gamma_c$ is zero, and sum of $c^2\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class CBDQCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDQCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CBDS

CBD model

Description

Fits and forecasts mortality rates using CBD model.

Usage

```
CBDS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD (M5) model is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \epsilon_{x,t}.$$

The model is estimated by regression and is forecasted by ARIMA applied to $\kappa_{1,t}$ and $\kappa_{2,t}$. It is designed for ages 50-90.

Value

An object of class CBDS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Cairns, A.J.G., Blake, D., and Dowd, K. (2006). A two-factor model for stochastic mortality with parameter uncertainty: Theory and calibration. *Journal of Risk and Insurance*, 73(4), 687-718.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
fit <- CBDS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CC

Mortality curve calculation

Description

Calculates mortality rates given mortality curve parameter values.

Usage

```
CC(
  x,
  par,
  curve = c("gompertz", "makeham", "oppermann", "thiele", "wittsteinbumsted", "perks",
    "weibull", "vandermaen", "beard", "heligmanpollard", "rogersplanck", "siler",
    "martinelle", "thatcher")
)
```

Arguments

x	vector of ages.
par	vector of mortality curve parameter values.
curve	name of mortality curve (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher).

Details

See MC() for more details of different mortality curves.

Value

Calculated mortality rates based on selected mortality curve and parameter values.

Examples

```
CC(x=60:89,par=c(0.0000082,0.10771),curve="gompertz")
```

CFM2S

Common factor model with global age pattern

Description

Fits and forecasts mortality rates of two populations using common factor model with global age pattern.

Usage

```
CFM2S(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model with global age pattern is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_x \kappa_{t,i} + \epsilon_{x,t,i}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of β_x is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CFM2S with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J., Wang, M., Liu, J., and Leung, J.W.Y. (2026). Financial valuation of retirement village via stochastic modelling of disability prevalence rates. *ASTIN Bulletin*, 56(2), 447-473.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b <- c(-0.00010,0.01195,0.03030,0.02170,0.03690,0.02365,0.02280,0.03850,0.05845,0.04415,
0.04185,0.05175,0.03670,0.04195,0.04090,0.02775,0.04990,0.02865,0.03935,0.03820,
0.04000,0.02790,0.03705,0.03370,0.02940,0.02850,0.03400,0.02310,0.02675,0.03430)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CFM2S(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

CFMS

Common factor model

Description

Fits and forecasts mortality rates of two populations using common factor model.

Usage

```
CFMS(x, M1, M2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M1	matrix of mortality rates of population 1 (rows as years and columns as ages).
M2	matrix of mortality rates of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model (CFM) is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_{x,i} \kappa_{t,i} + \epsilon_{x,t,i}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of $\beta_{x,i}$ is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class CFMS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, N. and Lee, R. (2005). Coherent mortality forecasts for a group of populations: An extension of the Lee-Carter method. *Demography*, 42(3), 575-594.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
```

```

2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b1 <- c(0.0012,-0.0033,0.0523,0.0161,0.0529,0.0220,0.0312,0.0437,0.0709,0.0444,
0.0398,0.0361,0.0403,0.0396,0.0506,0.0315,0.0428,0.0261,0.0384,0.0388,
0.0300,0.0269,0.0275,0.0256,0.0239,0.0421,0.0314,0.0284,0.0174,0.0314)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
b2 <- c(-0.0014,0.0272,0.0083,0.0273,0.0209,0.0253,0.0144,0.0333,0.0460,0.0439,
0.0439,0.0674,0.0331,0.0443,0.0312,0.0240,0.0570,0.0312,0.0403,0.0376,
0.0500,0.0289,0.0466,0.0418,0.0349,0.0149,0.0366,0.0178,0.0361,0.0372)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b1)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b2)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
fit <- CFMS(x=x,M1=M1,M2=M2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

curveS3

*Common S3 methods for mortality curves***Description**

Common S3 methods for objects returned by MC():

coef(object) gives estimated parameter values.

fitted(object) gives fitted values.

predict(object,x) gives estimated mortality rate(s) for given age(s) x.

plot(object) plots data and fitted mortality curve.

deviance(object) gives weighted sum of squared residuals on $\ln(m_x)$.

residuals(object) gives residuals on $\ln(m_x)$.

ENI

*Mortality ensemble interval***Description**

Generates ensemble interval forecast of mortality rates.

Usage

```
ENI(..., width = 0.95, method = 1, wm, nsim = 10, seed = 123)
```

Arguments

<code>...</code>	fitted model objects returned by <code>LCS()</code> , <code>RHS()</code> , <code>APCS()</code> , <code>CBDS()</code> , <code>CBDSC()</code> , <code>CBDQCS()</code> , and / or <code>STARS()</code> .
<code>width</code>	coverage probability of interval (default = 0.95).
<code>method</code>	if 1, simple averaging; if 2, weighted averaging; if 3, envelope; if 4, interior trimming (default = 1).
<code>wm</code>	vector of weights for LC, RH, APC, M5, M6, M7, and / or STAR models if method = 2 (default = equal weights).
<code>nsim</code>	number of simulations (default = 10).
<code>seed</code>	seed for random number generator (default = 123).

Details

Ensemble interval forecast is constructed by combining interval forecasts from individual stochastic mortality models using different methods including simple averaging, weighted averaging, envelope, and interior trimming. See `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDSC()`, `CBDQCS()`, and `STARS()` for more details of different stochastic mortality models.

Value

An object of class ENI with associated S3 methods `forecast` and `plot`.

References

Li, J., Wang, M., Liu, J., and Tickle, L. (2025). Ensemble interval forecasts of mortality. *Scandinavian Actuarial Journal*, 2025(6), 598-616.

Examples

```
x <- 60:69
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962)
b <- c(0.0801, 0.0909, 0.0948, 0.0951, 0.0965, 0.1014, 0.1042, 0.1141, 0.1110, 0.1118)
k <- c(12.11, 10.69, 11.18, 9.64, 9.35, 8.21, 6.89, 5.74, 4.56, 3.60,
3.27, 2.04, 1.11, -0.44, -1.05, -1.03, -1.84, -2.90, -4.03, -4.12,
-5.18, -5.64, -6.00, -6.51, -6.91, -6.90, -8.32, -8.53, -9.69, -9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=10,byrow=TRUE)+rnorm(300,0,0.035))
fit1 <- LCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit2 <- RHS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit3 <- APCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit4 <- CBDS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
fit <- ENI(fit1,fit2,fit3,fit4)
forecast::forecast(fit)
plot(fit)
```

ENS

*Mortality ensemble***Description**

Fits and forecasts mortality rates using mortality ensemble.

Usage

```
ENS(x, M, wm = rep(1/7, 7), curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
wm	vector of weights for LC, RH, APC, M5, M6, M7, and STAR models (default = 1/7).
curve	name of mortality curve for smoothing ensemble mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

Ensemble forecast is obtained as a weighted average of forecasts from individual stochastic mortality models. See `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDCS()`, `CBDQCS()`, and `STARS()` for more details of different stochastic mortality models.

Value

An object of class ENS with associated S3 methods `forecast` and `plot`.

References

Li, J. (2023). A model stacking approach for forecasting mortality. *North American Actuarial Journal*, 27(3), 530-545.

Examples

```
x <- 60:69
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962)
b <- c(0.0801,0.0909,0.0948,0.0951,0.0965,0.1014,0.1042,0.1141,0.1110,0.1118)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.60,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.90,-4.03,-4.12,
-5.18,-5.64,-6.00,-6.51,-6.91,-6.90,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=10,byrow=TRUE)+rnorm(300,0,0.035))
fit <- ENS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
forecast::forecast(fit)
plot(fit)
```

FCS

Mortality model fitting

Description

Fits and forecasts mortality rates using different stochastic mortality models with different estimation methods.

Usage

```
FCS(
  x,
  M,
  model = c("LC", "RH", "APC", "CBD", "CBDC", "CBDQC", "STAR"),
  curve = NULL,
  h = 10,
  jumpoff = 1
)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
model	name of stochastic mortality model (including LC, RH, APC, CBD, CBDC, CBDQC, and STAR).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).

<code>h</code>	forecast horizon (default = 10).
<code>jumpoff</code>	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

See `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDCS()`, `CBDQCS()`, and `STARS()` for more details of different stochastic mortality models.

Value

An object of class based on selected stochastic mortality model with associated S3 methods `coef`, `forecast`, `plot`, and `residuals`.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
0.0396, 0.0399, 0.0397, 0.0386, 0.039, 0.0375, 0.0367, 0.0368, 0.035, 0.0354,
0.0336, 0.0323, 0.0313, 0.0295, 0.0282, 0.0265, 0.024, 0.0226, 0.0219, 0.0183)
k <- c(12.11, 10.69, 11.18, 9.64, 9.35, 8.21, 6.89, 5.74, 4.56, 3.6,
3.27, 2.04, 1.11, -0.44, -1.05, -1.03, -1.84, -2.9, -4.03, -4.12,
-5.18, -5.64, -6, -6.51, -6.91, -6.9, -8.32, -8.53, -9.69, -9.31)
set.seed(123)
M <- exp(outer(k, b) + matrix(a, nrow=30, ncol=30, byrow=TRUE) + rnorm(900, 0, 0.035))
fit <- FCS(x=x, M=M, model="LC", curve="makeham", h=30, jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

LCLS

Lee-Carter model for limited data

Description

Fits and forecasts mortality rates using Lee-Carter model with sparse data in irregular years.

Usage

```
LCLS(x, t, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
t	vector of years.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model is specified as

$$\ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \epsilon_{x,t}.$$

The model is estimated by singular value decomposition and is forecasted by random walk with drift applied to κ_t . Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class LCLS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Li, N., Lee, R., and Tuljapurkar, S. (2004). Using the Lee-Carter method to forecast mortality for populations with limited data. *International Statistical Review*, 72(1), 19-36.

Examples

```
x <- 60:89
t <- c(1991,1996,2001,2006,2011:2020)
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821,-1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,8.21,
3.27,-1.03,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
```

```

set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=14,ncol=30,byrow=TRUE)+rnorm(420,0,0.035))
fit <- LCLS(x=x,t=t,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

LCS

Lee-Carter model

Description

Fits and forecasts mortality rates using Lee-Carter model.

Usage

```
LCS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model is specified as

$$\ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \epsilon_{x,t}.$$

The model is estimated by singular value decomposition and is forecasted by ARIMA applied to κ_t . Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class LCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Lee, R.D. and Carter, L.R. (1992). Modeling and forecasting U.S. mortality. *Journal of the American Statistical Association*, 87(419), 659-671.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
0.0396, 0.0399, 0.0397, 0.0386, 0.039, 0.0375, 0.0367, 0.0368, 0.035, 0.0354,
0.0336, 0.0323, 0.0313, 0.0295, 0.0282, 0.0265, 0.024, 0.0226, 0.0219, 0.0183)
k <- c(12.11, 10.69, 11.18, 9.64, 9.35, 8.21, 6.89, 5.74, 4.56, 3.6,
3.27, 2.04, 1.11, -0.44, -1.05, -1.03, -1.84, -2.9, -4.03, -4.12,
-5.18, -5.64, -6, -6.51, -6.91, -6.9, -8.32, -8.53, -9.69, -9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- LCS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

MC

Mortality curve fitting

Description

Fits parametric mortality curves to observed mortality rates using multiple optimisation strategies.

Usage

```
MC(x, m, curve = .curves, w = rep(1, length(x)))
```

Arguments

x	vector of ages.
m	vector of mortality rates.
curve	name of mortality curve (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive).
w	vector of weights (default = 1).

Details

User can choose one of the following 14 mortality curves (with their suitable age ranges in brackets):

- Gompertz (1825) (ages 30-90) - $m_x = Be^{Cx}$
- Makeham (1860) (ages 20-90) - $m_x = A + Be^{Cx}$
- Oppermann (1870) (ages 0-20) - $m_x = \frac{A}{\sqrt{x+1}} + B + C\sqrt{x+1}$
- Thiele (1871) (ages 0-90) - $m_x = A_1e^{-B_1x} + A_2e^{-0.5B_2(x-C)^2} + A_3e^{B_3x}$
- Wittstein & Bumsted (1883) (ages 0-90) - $m_x = \frac{A^{-(Bx)^N}}{B} + A^{-(M-x)^N}$
- Perks (1932) (ages 20-100+) - $m_x = \frac{A+BC^x}{1+DC^x}$
- Weibull (1939) (ages 50-100+) - $m_x = Bx^C$
- Van der Maen (1943) (ages 80-100+) - $m_x = A + Bx + Cx^2 + \frac{I}{N-x}$
- Beard (1971) (ages 50-100+) - $m_x = \frac{Ae^{Cx}}{1+Be^{Cx}}$
- Heligman & Pollard (1980) (ages 0-100+) - $m_x = A^{(x+B)^C} + De^{-E(\ln(x)-\ln(F))^2} + \frac{GH^x}{1+GH^x}$
- Rogers & Planck (1983) (ages 0-100+) - $m_x = A_0 + A_1e^{-Ax} + A_2e^{-B(x-U)-e^{-C(x-U)}} + A_3e^{Dx}$
- Siler (1983) (ages 0-90) - $m_x = A_1e^{-B_1x} + A_2 + A_3e^{B_3x}$
- Martinelle (1987) (ages 20-100+) - $m_x = \frac{A+Be^{Cx}}{1+De^{Cx}} + Ee^{Cx}$
- Thatcher (1999) (ages 20-100+) - $m_x = A + \frac{Be^{Cx}}{1+De^{Cx}}$

The mortality curves are fitted by employing multiple optimisation strategies simultaneously (including PORT routines, Nelder-Mead method, and Levenberg-Marquardt algorithm) and selecting one with smallest weighted least squares on $\ln(m_x)$. Constrained parameterisations offer traditional interpretability while unconstrained parameterisations provide increased flexibility.

Value

An object of class based on selected mortality curve with associated S3 methods coef, fitted, predict, plot, deviance, and residuals.

References

- Gompertz, B. (1825). On the nature of the function expressive of the law of human mortality, and on a new mode of determining the value of life contingencies. Philosophical Transactions of the Royal Society of London, 115(1825), 513-583.
- Makeham, W.M. (1860). On the law of mortality and the construction of annuity tables. Journal of the Institute of Actuaries, 8(6), 301-310.
- Oppermann, L.H.F. (1870). On the graduation of life tables, with special application to the rate of mortality in infancy and childhood. The Institute of Actuaries.
- Thiele, T.N. (1871). On a mathematical formula to express the rate of mortality throughout the whole of life, tested by a series of observations made use of by the Danish Life Insurance Company of 1871. Journal of the Institute of Actuaries and Assurance Magazine, 16(5), 313-329.

- Wittstein, T. and Bumsted, D.A. (1883). The mathematical law of mortality. *Journal of the Institute of Actuaries and Assurance Magazine*, 24(3), 153-173.
- Perks, W. (1932). On some experiments in the graduation of mortality statistics. *Journal of the Institute of Actuaries*, 63(1), 12-57.
- Weibull, W. (1951). A statistical distribution function of wide applicability. *Journal of Applied Mechanics*, 18(3), 293-297.
- Beard, R.E. (1971). Some aspects of theories of mortality, cause of death analysis, forecasting and stochastic processes. *Biological Aspects of Demography*, 57-68.
- Heligman, L. and Pollard, J.H. (1980). The age pattern of mortality. *Journal of the Institute of Actuaries*, 107(1), 49-80.
- Rogers, A. and Planck, F. (1983). Model: A general program for estimating parametrized model schedules of fertility, mortality, migration, and marital and labor force status transitions. IIASA Working Paper WP-83-102.
- Siler, W. (1983). Parameters of mortality in human populations with widely varying life spans. *Statistics in Medicine*, 2(3), 373-380.
- Martinelle, S. (1987). A generalized Perks formula for old-age mortality. *Statistiska Centralbyran*.
- Thatcher, A.R. (1999). The long-term pattern of adult mortality and the highest attained age. *Journal of the Royal Statistical Society Series A*, 162(1), 5-43.
- Tabeau, E. (2001). A review of demographic forecasting models for mortality. *Forecasting Mortality in Developed Countries. European Studies of Population*, Vol 9.

Examples

```
x <- 60:89
set.seed(123); m <- 0.0000082*exp(0.10771*c(60:89)+rnorm(30,0,0.1))
fit <- MC(x=x,m=m,curve="gompertz")
coef(fit)
fitted(fit)
predict(fit,62.5)
plot(fit)
deviance(fit)
residuals(fit)
```

modelS3

Common S3 methods for stochastic mortality models

Description

Common S3 methods for objects returned by `LCS()`, `RHS()`, `APCS()`, `CBDS()`, `CBDSC()`, `CBDQCS()`, `STARS()`, `ENS()`, `ENI()`, `CFMS()`, `CFM2S()`, `CAES()`, `PLCS()`, `PRHS()`, `PAPCS()`, `PCBDS()`, `PCBDSC()`, `PCBDQCS()`, `PCFMS()`, `PCFM2S()`, `PCAES()`, and `LCLS()`.

`coef(object)` gives estimated parameter values.

`forecast::forecast(object)` gives mortality forecasts (which = 1 for smoothed (default); which = 2 for raw) (or ensemble interval forecasts for `ENI()`).

plot(object) plots estimated parameter values, standardised residuals heatmap, data, and mortality forecasts smoothed by selected mortality curve (or ensemble interval forecasts for ENI()).

residuals(object) gives standardised residuals on $\ln(m_{x,t})$ or $D_{x,t}$.

simulate(object,nsim,seed) gives simulated future mortality rates (raw).

mortalitymatrix	<i>Convert mortality data to matrix</i>
-----------------	---

Description

Converts mortality rates stored as a matrix, data frame, or demogdata object into required matrix format.

Usage

```
mortalitymatrix(data, series = 1)
```

Arguments

data	mortality data.
series	mortality series index for demogdata objects.

Value

A numeric matrix of mortality data.

Examples

```
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
Data <- as.data.frame(M)
mortalitymatrix(Data)
```

PAPCS

*Poisson age-period-cohort model***Description**

Fits and forecasts mortality rates using age-period-cohort model with Poisson assumption.

Usage

```
PAPCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The age-period-cohort (APC) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \alpha_x + \kappa_t + \gamma_{t-x} \text{ and } D_{x,t} \text{ Poisson}(E_{x,t}m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c . Constraints include sum of κ_t is zero and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class PAPCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821,-1.7697)
b <- rep(1/30,30)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PAPCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCAES

Poisson common age effect model

Description

Fits and forecasts mortality rates of two populations using common age effect model with Poisson assumption.

Usage

```
PCAES(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2,

vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).

h forecast horizon (default = 10).

jumpoff if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common age effect (CAE) model with Poisson assumption is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + \beta_x \kappa_{t,i} \text{ and } D_{x,t,i} \text{ Poisson}(E_{x,t,i} m_{x,t,i}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{t,i}$. Constraints include sum of β_x is one and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCAES with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Kleinow, T. (2015). A common age effect model for the mortality of multiple populations. Insurance: Mathematics and Economics, 63(C), 147-152.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
b <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
k1 <- c(8.68,8.34,7.99,6.87,8.18,5.73,4.83,5.20,2.74,3.22,
2.99,1.59,1.67,-0.65,-0.39,-1.07,-0.95,-2.78,-3.46,-2.45,
-4.12,-4.66,-4.98,-4.58,-6.30,-4.39,-5.56,-6.52,-8.26,-6.92)
k2 <- c(11.81,11.01,10.59,10.40,9.75,8.15,6.07,6.45,4.60,4.57,
4.15,1.49,1.77,-1.08,-1.44,-0.96,-1.66,-2.25,-4.67,-4.62,
-4.38,-6.37,-6.27,-6.91,-8.22,-7.35,-8.39,-7.87,-9.72,-8.65)
set.seed(123)
M1 <- exp(outer(k1,b)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
```

```

E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCAES(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

PCBDCS

Poisson CBD with cohort model

Description

Fits and forecasts mortality rates using CBD with cohort model with Poisson assumption.

Usage

```
PCBDCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with cohort (M6) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \gamma_{t-x} \text{ and } D_{x,t} \text{ Poisson}(E_{x,t}m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, and γ_c . Constraints include sum of γ_c is zero and sum of $c\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class PCBDCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. *North American Actuarial Journal*, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDQCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCBDQCS

Poisson CBD with curvature and cohort model

Description

Fits and forecasts mortality rates using CBD with curvature and cohort model with Poisson assumption.

Usage

```
PCBDQCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD with curvature and cohort (M7) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) + \kappa_{3,t}((x - \bar{x})^2 - \sigma^2) + \gamma_{t-x} \text{ and } D_{x,t} \text{ Poisson}(E_{x,t}m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$, $\kappa_{2,t}$, $\kappa_{3,t}$, and γ_c . Constraints include sum of γ_c is zero, sum of $c\gamma_c$ is zero, and sum of $c^2\gamma_c$ is zero. It is designed for ages 50-90.

Value

An object of class PCBDQCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., Dowd, K., Coughlan, G.D., Epstein, D., Ong, A., and Balevich, I. (2009). A quantitative comparison of stochastic mortality models using data from England and Wales and the United States. North American Actuarial Journal, 13(1), 1-35.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDQCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCBDS

Poisson CBD model

Description

Fits and forecasts mortality rates using CBD model with Poisson assumption.

Usage

```
PCBDS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, perks, weibull, beard, martinelle, thatcher, gompertz2, makeham2, perks2, weibull2, beard2, martinelle2, thatcher2, where first 7 curves' parameters are unconstrained and last 7 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The CBD (M5) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \kappa_{1,t} + \kappa_{2,t}(x - \bar{x}) \text{ and } D_{x,t} \text{ Poisson}(E_{x,t}m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to $\kappa_{1,t}$ and $\kappa_{2,t}$. It is designed for ages 50-90.

Value

An object of class PCBDS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Cairns, A.J.G., Blake, D., and Dowd, K. (2006). A two-factor model for stochastic mortality with parameter uncertainty: Theory and calibration. *Journal of Risk and Insurance*, 73(4), 687-718.

Examples

```
x <- 60:89
k1 <- -2.97-0.0245*(0:29)
k2 <- 0.101+0.000345*(0:29)
set.seed(123)
M <- exp(matrix(k1,nrow=30,ncol=30,byrow=FALSE)+outer(k2,(x-mean(x)))+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PCBDS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

PCFM2S

*Poisson common factor model with global age pattern***Description**

Fits and forecasts mortality rates of two populations using common factor model with global age pattern with Poisson assumption.

Usage

```
PCFM2S(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model with global age pattern with Poisson assumption is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_x \kappa_{t,i} \text{ and } D_{x,t,i} \text{ Poisson}(E_{x,t,i} m_{x,t,i}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of β_x is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCFM2S with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J., Wang, M., Liu, J., and Leung, J.W.Y. (2026). Financial valuation of retirement village via stochastic modelling of disability prevalence rates. *ASTIN Bulletin*, 56(2), 447-473.

Examples

```
x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b <- c(-0.00010,0.01195,0.03030,0.02170,0.03690,0.02365,0.02280,0.03850,0.05845,0.04415,
0.04185,0.05175,0.03670,0.04195,0.04090,0.02775,0.04990,0.02865,0.03935,0.03820,
0.04000,0.02790,0.03705,0.03370,0.02940,0.02850,0.03400,0.02310,0.02675,0.03430)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCFM2S(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

Description

Fits and forecasts mortality rates of two populations using common factor model with Poisson assumption.

Usage

```
PCFMS(x, D1, D2, E1, E2, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D1	matrix of death counts of population 1 (rows as years and columns as ages).
D2	matrix of death counts of population 2 (rows as years and columns as ages).
E1	matrix of mid-year exposures of population 1 (rows as years and columns as ages).
E2	matrix of mid-year exposures of population 2 (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The common factor model (CFM) with Poisson assumption is specified as

$$\ln(m_{x,t,i}) = \alpha_{x,i} + B_x K_t + \beta_{x,i} \kappa_{t,i} \text{ and } D_{x,t,i} \text{ Poisson}(E_{x,t,i} m_{x,t,i}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to K_t and $\kappa_{t,i}$. Constraints include sum of B_x is one, sum of K_t is zero, sum of $\beta_{x,i}$ is one, and sum of $\kappa_{t,i}$ is zero. It can be applied to whole age range.

Value

An object of class PCFMS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot (which = 1 gives parameter estimates (default); which = 2 gives residuals and forecasts), and residuals.

References

Li, J. (2013). A Poisson common factor model for projecting mortality and life expectancy jointly for females and males. *Population Studies*, 67(1), 111-126.

Examples

```

x <- 60:89
a1 <- c(-5.18,-5.12,-4.98,-4.92,-4.82,-4.73,-4.66,-4.53,-4.45,-4.35,
-4.26,-4.17,-4.05,-3.95,-3.84,-3.73,-3.65,-3.52,-3.40,-3.29,
-3.14,-3.02,-2.88,-2.76,-2.64,-2.49,-2.37,-2.25,-2.12,-2.00)
a2 <- c(-4.78,-4.68,-4.57,-4.49,-4.39,-4.29,-4.19,-4.10,-4.00,-3.89,
-3.80,-3.69,-3.60,-3.49,-3.39,-3.29,-3.17,-3.07,-2.96,-2.85,
-2.71,-2.62,-2.49,-2.37,-2.26,-2.14,-2.04,-1.91,-1.82,-1.72)
B <- c(0.0381,0.0340,0.0420,0.0389,0.0423,0.0414,0.0406,0.0393,0.0415,0.0400,
0.0411,0.0362,0.0387,0.0381,0.0384,0.0385,0.0356,0.0314,0.0317,0.0337,
0.0316,0.0298,0.0284,0.0270,0.0248,0.0262,0.0205,0.0215,0.0142,0.0145)
K <- c(9.66,9.89,10.66,9.83,9.52,7.39,7.64,6.36,2.32,4.18,
2.91,-0.61,0.28,-0.38,-1.79,-3.34,-1.74,-3.50,-4.28,-4.77,
-4.98,-7.13,-5.09,-6.41,-5.56,-5.65,-6.12,-5.64,-7.35,-6.28)
b1 <- c(0.0012,-0.0033,0.0523,0.0161,0.0529,0.0220,0.0312,0.0437,0.0709,0.0444,
0.0398,0.0361,0.0403,0.0396,0.0506,0.0315,0.0428,0.0261,0.0384,0.0388,
0.0300,0.0269,0.0275,0.0256,0.0239,0.0421,0.0314,0.0284,0.0174,0.0314)
k1 <- c(-1.24,-1.38,-3.48,-2.51,-1.32,-1.90,-3.42,-0.94,0.24,-0.48,
-0.26,2.70,1.39,-0.46,1.74,2.53,0.90,1.43,0.76,2.48,
0.74,2.32,0.42,1.69,-0.64,1.30,0.19,-0.69,-1.11,-1.01)
b2 <- c(-0.0014,0.0272,0.0083,0.0273,0.0209,0.0253,0.0144,0.0333,0.0460,0.0439,
0.0439,0.0674,0.0331,0.0443,0.0312,0.0240,0.0570,0.0312,0.0403,0.0376,
0.0500,0.0289,0.0466,0.0418,0.0349,0.0149,0.0366,0.0178,0.0361,0.0372)
k2 <- c(2.35,0.62,-0.38,0.12,0.00,0.80,-1.39,0.38,2.47,0.40,
0.76,3.06,1.42,-0.73,0.79,1.94,0.12,0.60,-0.43,0.29,
0.17,0.98,-1.01,-0.13,-2.46,-1.24,-1.65,-2.48,-2.32,-3.06)
set.seed(123)
M1 <- exp(outer(k1,b1)+outer(K,B)+matrix(a1,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
M2 <- exp(outer(k2,b2)+outer(K,B)+matrix(a2,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.07))
E1 <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
E2 <- E1
D1 <- round(E1*M1)
D2 <- round(E2*M2)
fit <- PCFMS(x=x,D1=D1,D2=D2,E1=E1,E2=E2,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

Description

Fits and forecasts mortality rates using different stochastic mortality models with Poisson assumption.

Usage

```
PFCS(
  x,
  D,
  E,
  model = c("LC", "RH", "APC", "CBD", "CBDC", "CBDQC"),
  curve = NULL,
  h = 10,
  jumpoff = 1
)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
model	name of stochastic mortality model (including LC, RH, APC, CBD, CBDC, and CBDQC).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

See `PLCS()`, `PRHS()`, `PAPCS()`, `PCBDS()`, `PCBDCS()`, and `PCBDQCS()` for more details of different stochastic mortality models with Poisson assumption.

Value

An object of class based on selected stochastic mortality model with associated S3 methods `coef`, `forecast`, `plot`, and `residuals`.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
```

```

0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PFCS(x=x,D=D,E=E,model="LC",curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

PLCS

Poisson Lee-Carter model

Description

Fits and forecasts mortality rates using Lee-Carter model with Poisson assumption.

Usage

```
PLCS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Lee-Carter (LC) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t \text{ and } D_{x,t} \sim \text{Poisson}(E_{x,t} m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t . Constraints include sum of β_x is one and sum of κ_t is zero. It can be applied to whole age range.

Value

An object of class PLCS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Renshaw, A.E. and Haberman, S. (2006). A cohort-based extension to the Lee-Carter model for mortality reduction factors. *Insurance: Mathematics and Economics*, 38(3), 556-570.

Examples

```
x <- 60:89
a <- c(-4.8499,-4.7676,-4.6719,-4.5722,-4.4847,-4.3841,-4.2813,-4.1863,-4.0861,-3.9962,
-3.8885,-3.7896,-3.6853,-3.5737,-3.4728,-3.3718,-3.2586,-3.1474,-3.0371,-2.9206,
-2.7998,-2.6845,-2.5653,-2.4581,-2.3367,-2.2159,-2.1017,-1.9941,-1.8821, -1.7697)
b <- c(0.0283,0.0321,0.0335,0.0336,0.0341,0.0358,0.0368,0.0403,0.0392,0.0395,
0.0396,0.0399,0.0397,0.0386,0.039,0.0375,0.0367,0.0368,0.035,0.0354,
0.0336,0.0323,0.0313,0.0295,0.0282,0.0265,0.024,0.0226,0.0219,0.0183)
k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PLCS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

Description

Fits and forecasts mortality rates using Renshaw-Haberman model with Poisson assumption.

Usage

```
PRHS(x, D, E, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
D	matrix of death counts (rows as years and columns as ages).
E	matrix of mid-year exposures (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Renshaw-Haberman (RH) model with Poisson assumption is specified as

$$\ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \gamma_{t-x} \text{ and } D_{x,t} \text{ Poisson}(E_{x,t} m_{x,t}).$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c . Constraints include sum of β_x is one, sum of κ_t is zero, and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class PRHS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, and residuals.

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
0.0396, 0.0399, 0.0397, 0.0386, 0.039, 0.0375, 0.0367, 0.0368, 0.035, 0.0354,
0.0336, 0.0323, 0.0313, 0.0295, 0.0282, 0.0265, 0.024, 0.0226, 0.0219, 0.0183)
```

```

k <- c(12.11,10.69,11.18,9.64,9.35,8.21,6.89,5.74,4.56,3.6,
3.27,2.04,1.11,-0.44,-1.05,-1.03,-1.84,-2.9,-4.03,-4.12,
-5.18,-5.64,-6,-6.51,-6.91,-6.9,-8.32,-8.53,-9.69,-9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
E <- matrix(c(107788,108036,107481,106552,104608,100104,95803,91345,84980,79557,
75146,70559,65972,60898,55623,50522,47430,45895,41443,34774,
30531,27754,25105,22271,19437,16888,14458,12146,10038,7994),30,30,byrow=TRUE)
D <- round(E*M)
fit <- PRHS(x=x,D=D,E=E,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)

```

RHS

Renshaw-Haberman model

Description

Fits and forecasts mortality rates using Renshaw-Haberman model.

Usage

```
RHS(x, M, curve = NULL, h = 10, jumpoff = 1)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).
jumpoff	if 1, forecasts are based on estimated parameters only; if 2, forecasts are anchored to observed mortality rates in final year (default = 1).

Details

The Renshaw-Haberman (RH) model is specified as

$$\ln(m_{x,t}) = \alpha_x + \beta_x \kappa_t + \gamma_{t-x} + \epsilon_{x,t}.$$

The model is estimated by Newton updating scheme and is forecasted by ARIMA applied to κ_t and γ_c . Constraints include sum of β_x is one, sum of κ_t is zero, and sum of γ_c is zero. It can be applied to whole age range.

Value

An object of class RHS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Haberman, S. and Renshaw, A. (2011). A comparative study of parametric mortality projection models. Insurance: Mathematics and Economics, 48(1), 35-55.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
0.0396, 0.0399, 0.0397, 0.0386, 0.039, 0.0375, 0.0367, 0.0368, 0.035, 0.0354,
0.0336, 0.0323, 0.0313, 0.0295, 0.0282, 0.0265, 0.024, 0.0226, 0.0219, 0.0183)
k <- c(12.11, 10.69, 11.18, 9.64, 9.35, 8.21, 6.89, 5.74, 4.56, 3.6,
3.27, 2.04, 1.11, -0.44, -1.05, -1.03, -1.84, -2.9, -4.03, -4.12,
-5.18, -5.64, -6, -6.51, -6.91, -6.9, -8.32, -8.53, -9.69, -9.31)
set.seed(123)
M <- exp(outer(k,b)+matrix(a,nrow=30,ncol=30,byrow=TRUE)+rnorm(900,0,0.035))
fit <- RHS(x=x,M=M,curve="makeham",h=30,jumpoff=2)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

STARS

Spatial-temporal autoregressive model

Description

Fits and forecasts mortality rates using spatial-temporal autoregressive model.

Usage

```
STARS(x, M, curve = NULL, h = 10)
```

Arguments

x	vector of ages.
M	matrix of mortality rates (rows as years and columns as ages).
curve	name of mortality curve for smoothing forecasted mortality rates (including gompertz, makeham, oppermann, thiele, wittsteinbumsted, perks, weibull, vandermaen, beard, heligmanpollard, rogersplanck, siler, martinelle, thatcher, gompertz2, makeham2, oppermann2, thiele2, wittsteinbumsted2, perks2, weibull2, vandermaen2, beard2, heligmanpollard2, rogersplanck2, siler2, martinelle2, thatcher2, where first 14 curves' parameters are unconstrained and last 14 curves' parameters are generally restricted to be positive) (optional; if missing, no smoothing is performed).
h	forecast horizon (default = 10).

Details

The spatial-temporal autoregressive (STAR) model is specified as

$$\ln(M_t) = \mu + R\ln(M_{t-1}) + E_t,$$

where μ contains intercept parameters and R is banded matrix with non-zero entries in main diagonal and two subdiagonals below that.

The model is estimated by constrained regression and is forecasted via its autoregressive nature. Constraints in R include: all non-zero entries are between zero and one and sum of each row is one. It can be applied to whole age range.

Value

An object of class STARS with associated S3 methods coef, forecast (which = 1 for smoothed (default); which = 2 for raw), plot, residuals, and simulate (nsim for setting number of simulations; seed for initialising random number generator).

References

Li, H. and Lu, Y. (2017). Coherent forecasting of mortality rates: A sparse vector-autoregression approach. ASTIN Bulletin, 47(2), 563-600.

Examples

```
x <- 60:89
a <- c(-4.8499, -4.7676, -4.6719, -4.5722, -4.4847, -4.3841, -4.2813, -4.1863, -4.0861, -3.9962,
-3.8885, -3.7896, -3.6853, -3.5737, -3.4728, -3.3718, -3.2586, -3.1474, -3.0371, -2.9206,
-2.7998, -2.6845, -2.5653, -2.4581, -2.3367, -2.2159, -2.1017, -1.9941, -1.8821, -1.7697)
b <- c(0.0283, 0.0321, 0.0335, 0.0336, 0.0341, 0.0358, 0.0368, 0.0403, 0.0392, 0.0395,
0.0396, 0.0399, 0.0397, 0.0386, 0.039, 0.0375, 0.0367, 0.0368, 0.035, 0.0354,
0.0336, 0.0323, 0.0313, 0.0295, 0.0282, 0.0265, 0.024, 0.0226, 0.0219, 0.0183)
k <- c(12.11, 10.69, 11.18, 9.64, 9.35, 8.21, 6.89, 5.74, 4.56, 3.6,
3.27, 2.04, 1.11, -0.44, -1.05, -1.03, -1.84, -2.9, -4.03, -4.12,
-5.18, -5.64, -6, -6.51, -6.91, -6.9, -8.32, -8.53, -9.69, -9.31)
set.seed(123)
M <- exp(outer(k, b) + matrix(a, nrow=30, ncol=30, byrow=TRUE) + rnorm(900, 0, 0.035))
```

```
fit <- STARS(x=x,M=M,curve="makeham",h=30)
coef(fit)
forecast::forecast(fit)
plot(fit)
residuals(fit)
```

Index

APCS, [2](#)

CAES, [4](#)

CBDACS, [5](#)

CBDQCS, [6](#)

CBDS, [8](#)

CC, [9](#)

CFM2S, [10](#)

CFMS, [11](#)

curveS3, [13](#)

ENI, [13](#)

ENS, [15](#)

FCS, [16](#)

LCLS, [17](#)

LCS, [19](#)

MC, [20](#)

modelS3, [22](#)

mortalitymatrix, [23](#)

PAPCS, [24](#)

PCAES, [25](#)

PCBDACS, [27](#)

PCBDQCS, [28](#)

PCBDS, [29](#)

PCFM2S, [31](#)

PCFMS, [32](#)

PFCS, [34](#)

PLCS, [36](#)

PRHS, [37](#)

RHS, [39](#)

STARS, [40](#)