

Package ‘normality’

August 22, 2026

Title Tests for Departure from Normality

Version 0.0.4

Description

A toolkit for assessing data normality using a comprehensive collection of statistical methods. It includes descriptive measures and formal hypothesis tests, such as skewness and kurtosis tests, the Anderson–Darling test, the Shapiro–Wilk test, and the D'Agostino–Pearson K2 omnibus test.

License MIT + file LICENSE

URL <https://github.com/P10911004-NPUST/normality>

BugReports <https://github.com/P10911004-NPUST/normality/issues>

Encoding UTF-8

Depends R (>= 3.5)

LazyData true

Suggests testthat (>= 3.0.0)

Config/testthat/edition 3

Config/roxygen2/version 8.1.0

NeedsCompilation no

Author Joon-Keat Lai [aut, cre, cph] (ORCID:
<<https://orcid.org/0000-0002-9840-5836>>)

Maintainer Joon-Keat Lai <p10911004@gmail.com>

Repository CRAN

Date/Publication 2026-08-22 17:40:02 UTC

Contents

Anderson_Darling_test	2
check_normality	3
Cramer_von_Mises_test	5
D.Agostino_Pearson_test	6
is_normal	7
is_tied	8

Jarque_Bera_test	8
kurtosis	9
Lilliefors_test	11
normality_standard_output	12
Shapiro_Wilk_coef_table	13
Shapiro_Wilk_pval_table	13
Shapiro_Wilk_test	14
skewness	15
Index	17

Anderson_Darling_test	<i>Anderson-Darling Normality Test</i>
-----------------------	--

Description

Performs the Anderson–Darling (A2) normality test, an EDF-based goodness-of-fit test that gives greater weight to deviations in the tails of the distribution.

Usage

```
Anderson_Darling_test(  
  x,  
  alpha = 0.05,  
  silent = FALSE,  
  summary = TRUE,  
  misc = FALSE  
)
```

Arguments

- x A numeric vector, at least length of 8.
- alpha Numeric (default: 0.05). Significance threshold, range from 0 to 1.
- silent Logical (default: FALSE). If FALSE, print out the results.
- summary Logical (default: TRUE). Produce a summary table.
- misc Logical (default: FALSE). Output other unimportant parameters.

Value

A list.

References

- D'Agostino, R.B., 2017. Tests for the Normal Distribution. In: D'Agostino, R.B., Stephens, M.A. (Eds.), Goodness-of-Fit Techniques, 1st ed. Routledge, New York, pg. 372–373. <https://doi.org/10.1201/9780203753064>
- Stephens, M.A., 2017. Tests Based on EDF Statistics. In: D'Agostino, R.B., Stephens, M.A. (Eds.), Goodness-of-Fit Techniques, 1st ed. Routledge, New York, (Table 4.8 & 4.9) pg. 126–128. <https://doi.org/10.1201/9780203753064>
- Anderson, T.W., Darling, D.A., 1954. A Test of Goodness of Fit. J. Am. Stat. Assoc. 49, 765–769. <https://doi.org/10.1080/01621459.1954.10501232>

Examples

```
out <- Anderson_Darling_test(rnorm(10))
print(out$summary)
```

check_normality	<i>Normality test</i>
-----------------	-----------------------

Description

A wrapper function for the normality tests available in this package.

Usage

```
check_normality(
  data,
  formula = NULL,
  alpha = 0.05,
  method = "SWR",
  summary = TRUE
)
```

Arguments

data	A data frame or a numeric vector.
formula	Formula (default: NULL). If data is a data frame, define the val ~ group.
alpha	Numeric (default: 0.05). Significance level used to determine whether the null hypothesis is rejected. Must be between 0 and 1.
method	Character. Abbreviation specifying the normality test to perform. Available options are c("AD", "CVM", "DAP", "JB", "LF", "SW", "SF", "SWR").
summary	Logical (default: TRUE). If TRUE, return a summary table of the test results.

Details

The method argument specifies the statistical procedure used to assess whether a sample is consistent with a normal distribution. Different tests emphasize different characteristics of departures from normality, such as skewness, kurtosis, or discrepancies in the tails of the distribution. Because no single test performs optimally under all circumstances, the choice of method may depend on sample size and the expected type of non-normality.

Available methods are:

- "AD": Anderson–Darling test. A modification of the empirical distribution function (EDF) approach that gives greater weight to observations in the tails of the distribution. Compared with several alternative normality tests, the Anderson–Darling procedure is often more sensitive to deviations occurring in extreme values and tail behavior. This test is applicable only for sample sizes $n \geq 8$.
- "CVM": Cramér–von Mises test. An empirical distribution function (EDF)-based goodness-of-fit test that measures the overall discrepancy between the empirical and theoretical cumulative distribution functions by assigning relatively uniform weight across the entire distribution. Compared with the Anderson–Darling test, the Cramér–von Mises test is generally less sensitive to deviations in the tails but performs well for detecting overall departures from normality. This test is applicable only for sample sizes $n \geq 8$.
- "DAP": D’Agostino–Pearson test. A combined omnibus moment test based on sample skewness and kurtosis. The procedure transforms the skewness and kurtosis statistics into approximately standard normal variables and combines them into a single test statistic. This method is designed to detect a broad range of departures from normality rather than emphasizing any particular feature. This test is applicable only for sample sizes $n \geq 20$.
- "JB": Jarque–Bera test. An omnibus moment test based on sample skewness and kurtosis. The test evaluates whether the observed skewness and kurtosis differ significantly from the values expected under a normal distribution. The method is commonly used in econometrics and is generally more appropriate for moderate to large sample sizes.
- "LF": Lilliefors test. The Lilliefors test is an EDF omnibus test modified from Kolmogorov–Smirnov test for the composite hypothesis of normality. The test statistic is the maximal absolute difference between empirical and hypothetical cumulative distribution function.
- "SW": Shapiro–Wilk test. The original normality test proposed by Shapiro and Wilk (1965), based on the correlation between ordered observations and their expected values under normality. It is widely regarded as one of the most powerful tests for detecting departures from normality in small samples. Applicable only for sample sizes $3 \leq n \leq 50$.
- "SF": Shapiro–Francia test. Proposed by Shapiro and Francia (1972) and subsequently simplified and extended by Royston (1993). This method is a computationally simpler modification of the Shapiro–Wilk procedure that performs particularly well for detecting departures associated with heavier-tailed distributions. Applicable only for sample sizes $5 \leq n \leq 5000$.
- "SWR": Shapiro–Wilk test with Royston’s modifications. Uses Royston’s (1992) approximations for the null distribution of the Shapiro–Wilk statistic and extends applicability to larger samples while maintaining behavior similar to the original test. Applicable only for sample sizes $3 \leq n \leq 5000$.

In all methods, the null hypothesis is that the sample is drawn from a normal distribution. Small p-values indicate evidence against the assumption of normality.

Value

A list.

Examples

```
out_AD <- check_normality(rnorm(20), method = "AD")
out_DAP <- check_normality(rnorm(20), method = "DAP")
out_SW <- check_normality(rnorm(20), method = "SW")
```

Cramer_von_Mises_test *Cramer-von Mises Normality Test*

Description

An empirical distribution function (EDF)-based goodness-of-fit test that measures the overall discrepancy between the empirical and theoretical cumulative distribution functions by assigning relatively uniform weight across the entire distribution.

Usage

```
Cramer_von_Mises_test(
  x,
  alpha = 0.05,
  silent = FALSE,
  summary = TRUE,
  misc = FALSE
)
```

Arguments

x	A numeric vector, at least length of 8.
alpha	Numeric (default: 0.05). Significance threshold, range from 0 to 1.
silent	Logical (default: FALSE). If FALSE, print out the results.
summary	Logical (default: TRUE). Produce a summary table.
misc	Logical (default: FALSE). Output other unimportant parameters.

Value

A list.

References

Thode, H. C., Jr., 2002. Goodness of fit tests. Testing for normality. Marcel Dekker, New York. (Section 5.1.3, pg. 103)

Stephens, M.A., 2017. Tests Based on EDF Statistics. In: D'Agostino, R.B., Stephens, M.A. (Eds.), Goodness-of-Fit Techniques, 1st ed. Routledge, New York, (Table 4.9, pg. 127). <https://doi.org/10.1201/9780203753064>

Examples

```
out <- Cramer_von_Mises_test(rnorm(10))
print(out$summary)
```

D.Agostino_Pearson_test

D'Agostino–Pearson K2 Normality Test

Description

The D'Agostino–Pearson chi-squared (K2) test is a moment-based omnibus test for normality.

Usage

```
D.Agostino_Pearson_test(
  x,
  alpha = 0.05,
  alternative = c("two.sided", "less", "greater"),
  silent = FALSE,
  summary = TRUE,
  misc = FALSE
)
```

Arguments

x	Numeric vector. Must have length at least 20.
alpha	Numeric (default: 0.05). Significance level for hypothesis testing. Must be between 0 and 1.
alternative	Character (default: "two.sided"). Specifies the alternative hypothesis. Available options are c("two.sided", "less", "greater"). Note that this option is only applied to the skewness and kurtosis components of the test.
silent	Logical (default: FALSE). If FALSE, results are printed to the console.
summary	Logical (default: TRUE). Produce a summary table.
misc	Logical (default: FALSE). Output other unimportant parameters.

Details

It evaluates the null hypothesis that the data come from a normal distribution by combining standardized measures of skewness and kurtosis into a single chi-squared test statistic.

Value

A list

References

D'Agostino, R.B., Belanger, A., D'Agostino, R.B., 1990. A Suggestion for Using Powerful and Informative Tests of Normality. Am. Stat. 44, 316–321. <https://doi.org/10.1080/00031305.1990.10475751>

Examples

```
out <- D.Agostino_Pearson_test(rnorm(50))
print(out$summary)
```

is_normal	<i>Normality test</i>
-----------	-----------------------

Description

A handy wrapper for data normality assessment using the Shapiro-Wilk-Royston, D'Agostino-Pearson, and Anderson-Darling tests.

Usage

```
is_normal(data, formula = NULL, alpha = 0.05, sensitivity = 2, summary = FALSE)
```

Arguments

data	A data frame or a numeric vector.
formula	Formula (default: NULL). If data is a data frame, define the val ~ group.
alpha	Significance threshold, range from 0 to 1 (default: 0.05).
sensitivity	Numeric, range from 1 to 3 (default: 2). The greater the value, the greater chance to consider as non-normal.
summary	Logical (default: FALSE). If TRUE, show the summary table.

Value

A boolean value (or a list if summary = TRUE).

Examples

```
is_normal(iris, Sepal.Length ~ Species)
```

is_tied	<i>Tied data</i>
---------	------------------

Description

Tied data

Usage

```
is_tied(x, ratio = 0.3, remove_NA = FALSE)
```

Arguments

x	A numeric vector
ratio	Numeric (default: 0.3). The ratio threshold of being considered as tied-data. The value range from 0 to 1.
remove_NA	Logical (default: TRUE). Whether or not to remove NAs.

Value

Logical

Examples

```
is_tied(c(1, 1, 2, 2, 2, 3, 4, 5))
#> TRUE
```

Jarque_Bera_test	<i>Jarque-Bera Normality Test</i>
------------------	-----------------------------------

Description

Performs the Jarque-Bera chi-squared test, a moment-based omnibus test for assessing normality.

Usage

```
Jarque_Bera_test(
  x,
  alpha = 0.05,
  alternative = c("two.sided", "less", "greater"),
  silent = FALSE,
  summary = TRUE
)
```

Arguments

<code>x</code>	Numeric vector. Must contain at least 20 observations.
<code>alpha</code>	Numeric (default: 0.05). Significance level for hypothesis testing. Must be between 0 and 1.
<code>alternative</code>	Character (default: "two.sided"). Specifies the alternative hypothesis. Available options are <code>c("two.sided", "less", "greater")</code> . This argument applies only to the skewness and kurtosis components and does not affect the Jarque-Bera omnibus test statistic itself.
<code>silent</code>	Logical (default: FALSE). If FALSE, results are printed to the console.
<code>summary</code>	Logical (default: TRUE). Produce a summary table.

Details

The test evaluates the null hypothesis that the data are drawn from a normal distribution by combining standardized measures of skewness and kurtosis into a single chi-squared test statistic.

Value

A list

References

Jarque, C.M., Bera, A.K., 1987. A Test for Normality of Observations and Regression Residuals. *Int. Stat. Rev.* 55, 163–172. <https://doi.org/10.2307/1403192>

See Also

[D.Agostino_Pearson_test\(\)](#)

Examples

```
out <- Jarque_Bera_test(rnorm(50))
print(out$summary)
```

kurtosis

Kurtosis test

Description

Performs a kurtosis test to assess whether a distribution deviates from normality in terms of tail heaviness.

Usage

```
kurtosis(
  x,
  alpha = 0.05,
  alternative = c("two.sided", "less", "greater"),
  method = c("G2", "b2", "g2"),
  silent = FALSE,
  summary = TRUE
)
```

Arguments

<code>x</code>	Numeric vector containing the input data.
<code>alpha</code>	Numeric (default: 0.05). Significance level for hypothesis testing. Must be between 0 and 1.
<code>alternative</code>	Character (default: "two.sided"). Specifies the alternative hypothesis. Available options are c("two.sided", "less", "greater").
<code>method</code>	Character (default: "G2"). Formula used to estimate kurtosis. Available options are c("G2", "b2", "g2"). The "g2" statistic is the classical sample kurtosis estimator, while "G2" and "b2" are bias-corrected versions of "g2".
<code>silent</code>	Logical (default: FALSE). If FALSE, results are printed to the console.
<code>summary</code>	Logical (default: TRUE). Produce a summary table.

Details

The test evaluates the null hypothesis that the population kurtosis is equal to 3, which is the kurtosis of a normal distribution. Values significantly different from 3 indicate deviations from normality, such as heavy-tailed or light-tailed behavior.

Value

A list

References

Joanes, D.N., Gill, C.A., 1998. Comparing measures of sample skewness and kurtosis. *J. R. Stat. Soc. D (The Statistician)* 47, 183–189. <https://doi.org/10.1111/1467-9884.00122>

Wright, D.B., Herrington, J.A., 2011. Problematic standard errors and confidence intervals for skewness and kurtosis. *Behav. Res. Methods* 43, 8–17. <https://doi.org/10.3758/s13428-010-0044-x>

Examples

```
x <- c(10:17, 12, 12, 13, 13, 13, 13, 13, 14, 14, 14, 15, 15)
kurtosis(x)
```

Lilliefors_test	<i>Lilliefors Normality Test</i>
-----------------	----------------------------------

Description

Performs the Lilliefors normality test, which is an empirical distribution function (EDF)-based goodness-of-fit test derived from the Kolmogorov–Smirnov test, using the approximation proposed by Molin and Abdi (1998).

Usage

```
Lilliefors_test(x, alpha = 0.05, silent = FALSE, summary = TRUE, misc = FALSE)
```

Arguments

x	A numeric vector, at least length of 8.
alpha	Numeric (default: 0.05). Significance threshold, range from 0 to 1.
silent	Logical (default: FALSE). If FALSE, print out the results.
summary	Logical (default: TRUE). Produce a summary table.
misc	Logical (default: FALSE). Output other unimportant parameters.

Value

A list.

References

Molin, P., Abdi, H., 1998. New tables and numerical approximation for the Kolmogorov-Smirnov/Lilliefors/Van Soest test of normality. Technical report, University of Bourgogne.

Examples

```
out <- Lilliefors_test(rnorm(10))
print(out$summary)
```

normality_standard_output
Standard output format

Description

The standard output format for normality package.

Usage

```
normality_standard_output(  
  method = "what test?",  
  is_normal = NA,  
  alpha = NA_real_,  
  alternative = c("two.sided", "less", "greater"),  
  summary = NULL,  
  statistic = NA_real_,  
  pvalue = NA_real_,  
  misc = NULL  
)
```

Arguments

method	Character. The name of the test.
is_normal	Logical. Is the input data normally distributed?
alpha	Numeric (default: 0.05). Significance threshold.
alternative	Character. The alternative hypothesis (H1) to test. Available options are c("two.sided", "less", "greater").
summary	Statistic summary, if any.
statistic	Numeric. The value used to calculate p-value.
pvalue	Numeric. The p-value of the test.
misc	List. Miscellaneous elements.

Value

A list.

`Shapiro_Wilk_coef_table`*Shapiro-Wilk normality test (coefficients)*

Description

Coefficients (ai) for the W test for normality.

Usage

`Shapiro_Wilk_coef_table`

Format

A data frame with 50 rows and 25 variables:

rownames is the sample size (n); colnames is the corresponding coefficients (ai).

References

Shapiro, S.S., Wilk, M.B., 1965. An Analysis of Variance Test for Normality (Complete Samples). *Biometrika* 52, 591–611. <https://doi.org/10.2307/2333709>

`Shapiro_Wilk_pval_table`*Shapiro-Wilk normality test (p-values)*

Description

The percentage points (critical values of W) of the W test for $n = 3(1)50$.

Usage

`Shapiro_Wilk_pval_table`

Format

A data frame with 50 rows and 10 variables:

rownames is the sample size (n); colnames is the corresponding p-values.

References

Shapiro, S.S., Wilk, M.B., 1965. An Analysis of Variance Test for Normality (Complete Samples). *Biometrika* 52, 591–611. <https://doi.org/10.2307/2333709>

Shapiro_Wilk_test	<i>Shapiro-Wilk Normality Test</i>
-------------------	------------------------------------

Description

Performs the Shapiro–Wilk normality test, which assesses whether a sample originates from a normally distributed population using a regression-based correlation method.

Usage

```
Shapiro_Wilk_test(
  x,
  alpha = 0.05,
  method = c("SWR", "SF", "SW"),
  silent = FALSE,
  summary = TRUE,
  misc = FALSE,
  resampling = TRUE
)
```

Arguments

x	A numeric vector.
alpha	Significance threshold (default: 0.05).
method	Character (default: "SWR"). Use which modification of the test? Available options are c("SWR", "SF", "SW").
silent	Logical (default: FALSE). If FALSE, print out the results.
summary	Logical (default: TRUE). Produce a summary table.
misc	Logical (default: FALSE). Output other unimportant parameters.
resampling	Logical (default: TRUE). If TRUE, unlock the sample size limitation of the test by using sample resampling method.

Details**method**

- "SW": Shapiro–Wilk test, the original normality test proposed by Shapiro and Wilk (1965). Applicable only for sample sizes $3 \leq n \leq 50$.
- "SF": Shapiro–Francia test, proposed by Shapiro and Francia (1972) and subsequently simplified and extended by Royston (1993). Applicable only for sample sizes $5 \leq n \leq 5000$.
- "SWR": Shapiro–Wilk test with Royston’s (1992) modifications for approximating the null distribution and extending the test to larger sample sizes. Applicable only for sample sizes $3 \leq n \leq 5000$.

Value

A list.

References

Shapiro, S.S., Wilk, M.B., 1965. An Analysis of Variance Test for Normality (Complete Samples). *Biometrika* 52, 591–611. <https://doi.org/10.2307/2333709>

Shapiro, S.S., Francia, R.S., 1972. An Approximate Analysis of Variance Test for Normality. *J. Am. Stat. Assoc.* 67, 215–216. <https://doi.org/10.1080/01621459.1972.10481232>

Royston, P., 1993. A pocket-calculator algorithm for the Shapiro–Francia test for non-normality: an application to medicine. *Stat. Med.* 12, 181–184. <https://doi.org/10.1002/sim.4780120209>

Royston, P., 1992. Approximating the Shapiro–Wilk W-test for non-normality. *Stat. Comput.* 2, 117–119. <https://doi.org/10.1007/BF01891203>

Examples

```
sw <- Shapiro_Wilk_test(rnorm(20), method = "SW")
print(sw$summary)
sf <- Shapiro_Wilk_test(rnorm(100) ^ 2, method = "SF")
print(sf$summary)
swr <- Shapiro_Wilk_test(rnorm(1e6), method = "SWR")
print(swr$summary)
```

skewness	<i>Skewness test</i>
----------	----------------------

Description

The test evaluates whether the population skewness is equal to zero. Under the null hypothesis, the data are assumed to originate from a symmetric distribution. Significant positive or negative skewness indicates asymmetry in the distribution and may suggest a departure from normality.

Usage

```
skewness(
  x,
  alpha = 0.05,
  alternative = c("two.sided", "less", "greater"),
  method = c("G1", "b1", "g1"),
  silent = FALSE,
  summary = TRUE
)
```

Arguments

x	Numeric vector containing the input data.
alpha	Numeric (default: 0.05). Significance level for hypothesis testing. Must be between 0 and 1.
alternative	Character (default: "two.sided"). Specifies the alternative hypothesis. Available options are c("two.sided", "less", "greater").
method	Character (default: "G1"). Formula used to estimate skewness. Available options are c("G1", "b1", "g1"). The "g1" statistic is the conventional moment-based sample skewness. The "G1" and "b1" statistics apply finite-sample corrections to reduce the bias of "g1".
silent	Logical (default: FALSE). If FALSE, the test results are printed to the console.
summary	Logical (default: TRUE). Produce a summary table.

Value

A list

References

- Joanes, D.N., Gill, C.A., 1998. Comparing measures of sample skewness and kurtosis. *J. R. Stat. Soc. D (The Statistician)* 47, 183–189. <https://doi.org/10.1111/1467-9884.00122>
- Wright, D.B., Herrington, J.A., 2011. Problematic standard errors and confidence intervals for skewness and kurtosis. *Behav. Res. Methods* 43, 8–17. <https://doi.org/10.3758/s13428-010-0044-x>

Examples

```
skewness(rnorm(30))
```

Index

* datasets

Shapiro_Wilk_coef_table, [13](#)

Shapiro_Wilk_pval_table, [13](#)

Anderson_Darling_test, [2](#)

check_normality, [3](#)

Cramer_von_Mises_test, [5](#)

D.Agostino_Pearson_test, [6](#)

D.Agostino_Pearson_test(), [9](#)

is_normal, [7](#)

is_tied, [8](#)

Jarque_Bera_test, [8](#)

kurtosis, [9](#)

Lilliefors_test, [11](#)

normality_standard_output, [12](#)

Shapiro_Wilk_coef_table, [13](#)

Shapiro_Wilk_pval_table, [13](#)

Shapiro_Wilk_test, [14](#)

skewness, [15](#)